

AMERICAN ACADEMY OF PEDIATRICS

Committee on Public Education

Children, Adolescents, and Television

ABSTRACT. This statement describes the possible negative health effects of television viewing on children and adolescents, such as violent or aggressive behavior, substance use, sexual activity, obesity, poor body image, and decreased school performance. In addition to the television ratings system and the v-chip (electronic device to block programming), media education is an effective approach to mitigating these potential problems. The American Academy of Pediatrics offers a list of recommendations on this issue for pediatricians and for parents, the federal government, and the entertainment industry.

ABBREVIATIONS. AAP, American Academy of Pediatrics; MTV, Music Television; E/I, educational/informational.

For the past 15 years, the American Academy of Pediatrics (AAP) has expressed its concerns about the amount of time children and adolescents spend viewing television and the content of what they view.¹ According to recent Nielsen Media Research data, the average child or adolescent watches an average of nearly 3 hours of television per day.² This figure does not include time spent watching videotapes or playing video games³ (a 1999 study found that children spend an average of 6 hours 32 minutes per day with various media combined).⁴ By the time the average person reaches age 70, he or she will have spent the equivalent of 7 to 10 years watching television.⁵ One recent study found that 32% of 2- to 7-year-olds and 65% of 8- to 18-year-olds have television sets in their bedrooms.⁴ Time spent with various media may displace other more active and meaningful pursuits, such as reading, exercising, or playing with friends.

Although there are potential benefits from viewing some television shows, such as the promotion of positive aspects of social behavior (eg, sharing, manners, and cooperation), many negative health effects also can result. Children and adolescents are particularly vulnerable to the messages conveyed through television, which influence their perceptions and behaviors.⁶ Many younger children cannot discriminate between what they see and what is real. Research has shown primary negative health effects on violence and aggressive behavior⁷⁻¹²; sexuality^{7,13-15}; academic performance¹⁶; body concept and self-im-

age¹⁷⁻¹⁹; nutrition, dieting, and obesity^{17,20,21}; and substance use and abuse patterns.⁷

In the scientific literature on media violence, the connection of media violence to real-life aggressive behavior and violence has been substantiated.⁸⁻¹² As much as 10% to 20% of real-life violence may be attributable to media violence.²² The recently completed 3-year National Television Violence Study found the following: 1) nearly two thirds of all programming contains violence; 2) children's shows contain the most violence; 3) portrayals of violence are usually glamorized; and 4) perpetrators often go unpunished.²³ A recent comprehensive analysis of music videos found that nearly one fourth of all Music Television (MTV) videos portray overt violence and depict weapon carrying.²⁴ Research has shown that even television news can traumatize children or lead to nightmares.²⁵ In a random survey of parents with children in kindergarten through sixth grade, 37% reported that their child had been frightened or upset by a television story in the preceding year.²⁶

According to a recent content analysis, mainstream television programming contains large numbers of references to cigarettes, alcohol, and illicit drugs.²⁷ One fourth of all MTV videos contain alcohol or tobacco use.²⁸ A longitudinal study found a positive correlation between television and music video viewing and alcohol consumption among teens.²⁹ Finally, content analyses show that children and teenagers continue to be bombarded with sexual imagery and innuendoes in programming and advertising.^{14,30,31} To date, there are no data available to substantiate the behavioral impact of this exposure.³¹

The new television ratings system and the v-chip are tools that can help protect children from potentially harmful content. All new television sets with screens measuring 13 inches or greater contain a v-chip that enables parents to program televisions to block out any shows that they deem inappropriate for their children.³² To block out television shows, parents must use the television ratings system, which has age and content descriptors for violence, sexual situations, suggestive dialogue, and adult language. Although the ratings system and the v-chip can assist parents, ongoing evaluation is necessary to ensure that these tools are as effective as possible.³³⁻³⁵ For example, the ratings should be applied uniformly and listed in television guides, newspapers, and journals so parents know what they mean.

Besides the v-chip, there are other means of protecting children from what is on television. Evidence

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

PEDIATRICS (ISSN 0031 4005). Copyright © 2001 by the American Academy of Pediatrics.

now shows that media education can help mitigate the harmful effects of media violence³⁶⁻⁴⁰ and alcohol advertising^{41,42} on children and adolescents. Media education programs have been included in the school curricula beginning in early elementary school in many states across the United States.⁴³

Furthermore, continued support of the Children's Television Act of 1990⁴⁴ and additional regulations made in 1996⁴⁵ will help to ensure the airing of television programs specifically designated for children. The act requires broadcasters to air educational and informational programming for children at least 3 hours per week and to limit the amount of advertising time allowed during children's programming. The shows must be labeled E/I (for educational and informational) on the television screen.

RECOMMENDATIONS

The following recommendations are given for pediatricians and other health care professionals:

1. Remain knowledgeable about the effects of television, including violent and aggressive behavior, obesity, poor body concept and self-image, substance use, and early sexual activity, by becoming involved in the AAP *Media Matters* campaign.⁴⁶ Educate patients and their parents about these effects.
2. Use the AAP *Media History* form⁴⁶ to help parents recognize the extent of their children's media consumption.
3. Work with local schools to implement comprehensive media-education programs that deal with important public health issues.³⁶
4. Serve as good role models by using television appropriately and by implementing reading programs using volunteer readers in waiting rooms and hospital inpatient units.
5. Become involved in the AAP's Media Resource Team (contact the Division of Public Education), and learn how to work effectively with writers, directors, and producers to make media more appropriate for children and adolescents. Contact networks and producers of television programs with concerns about the content of specific shows and episodes.
6. Ensure that appropriate entertainment options are available for hospitalized children and adolescents. Work with child life staff to assemble a screening committee that selects programs for closed circuit broadcast or a video library. Develop institution-specific, formal guidelines based on the established ratings system (which takes profanity, sex, and violence into account), and screen for content containing ethnic and sex role stereotyping. Considerations should also be made to avoid themes hospitalized children might find upsetting, and efforts should be made to enforce the ratings system in the hospital setting.
7. Support the Children's Television Act of 1990 and its 1996 rules by working to ensure that local television stations are in compliance with the act and by urging local newspapers to list ratings and E/I denotations of programs.

8. Monitor the television ratings system for appropriateness and advocate for substantive, content-based ratings in the future.

Pediatricians should recommend the following guidelines for parents:

1. Limit children's total media time (with entertainment media) to no more than 1 to 2 hours of quality programming per day.
2. Remove television sets from children's bedrooms.
3. Discourage television viewing for children younger than 2 years, and encourage more interactive activities that will promote proper brain development, such as talking, playing, singing, and reading together.
4. Monitor the shows children and adolescents are viewing. Most programs should be informational, educational, and nonviolent.
5. View television programs along with children, and discuss the content. Two recent surveys involving a total of nearly 1500 parents found that less than half of parents reported always watching television with their children.^{5,47}
6. Use controversial programming as a stepping-off point to initiate discussions about family values, violence, sex and sexuality, and drugs.
7. Use the videocassette recorder wisely to show or record high-quality, educational programming for children.
8. Support efforts to establish comprehensive media-education programs in schools.
9. Encourage alternative entertainment for children, including reading, athletics, hobbies, and creative play.

Pediatricians should lead efforts in their communities to do the following:

1. Form coalitions including libraries, religious organizations, and other community groups to broaden media education beyond the schools.
2. Organize activities promoting media education, such as letter-writing campaigns to local television stations to advocate for better programming for children, and developing local TV turnoff week projects.⁴⁸

Pediatricians should work with the Academy and local chapters to challenge the federal government to do the following:

1. Initiate legislation and rules that would ban alcohol advertising from television.
2. Fund ongoing annual research, such as the National Television Violence Study, and fund more research on the effects of television on children and adolescents, particularly in the area of sex and sexuality.
3. Assemble a *National Institutes of Health Comprehensive Report on Children, Adolescents, and Media* that would bring together all of the current relevant research.
4. Work with the US Department of Education to support the creation and implementation of media-education curricula for school children.

Pediatricians should work with the Academy and local chapters to challenge the entertainment industry to do the following:

1. Take responsibility for the programming it produces.
2. Adhere to the current television ratings system, and label programs conscientiously.
3. Collaborate with other public health advocates to convene a series of seminars with writers, directors, and producers to discuss ways to make media more appropriate for children and adolescents.
4. Produce more educational programming for children and adolescents, and ensure that the programming it produces is of higher quality, with less content that is gratuitously violent, sexually suggestive, or drug oriented.

COMMITTEE ON PUBLIC EDUCATION, 2000–2001

Miriam E. Bar-on, MD, Chairperson
Daniel D. Broughton, MD
Susan Buttross, MD
Suzanne Corrigan, MD
Alberto Gedissman, MD
M. Rosario González de Rivas, MD
Michael Rich, MD, MPH
Donald L. Shifrin, MD

LIAISONS

Michael Brody, MD
American Academy of Child and Adolescent Psychiatry
Brian Wilcox, PhD
American Psychological Association

CONSULTANTS

Marjorie Hogan, MD
H. James Holroyd, MD
Linda Reid, MD
S. Norman Sherry, MD
Victor Strasburger, MD

STAFF

Jennifer Stone

REFERENCES

1. American Academy of Pediatrics, Task Force on Children, and Television. Children, adolescents and television. *News and Comment*. December 1984;35:8
2. 1998 Report on Television. New York, NY: Nielsen Media Research; 1998.
3. Mares ML. Children's use of VCRs. *Ann Am Acad Pol Soc Science*. 1998;557:120–131
4. Roberts DF, Foehr UG, Rideout VJ, Brodie, M. *Kids and Media at the New Millennium: A Comprehensive National Analysis of Children's Media Use*. Menlo Park, CA: The Henry J Kaiser Family Foundation Report; 1999
5. Strasburger VC. Children, adolescents, and the media: five crucial issues. *Adolesc Med*. 1993;4:479–493
6. Gerbner G, Gross L, Morgan M, Signorelli N. Growing up with television: the cultivation perspective. In Bryant J, Zillmann D, eds. *Media Effects: Advances in Theory and Research*. Hillsdale, NJ: Lawrence Erlbaum; 1994:17–41
7. Strasburger VC. "Sex, drugs, rock'n'roll," and the media: are the media responsible for adolescent behavior? *Adolesc Med*. 1997;8:403–414
8. Strasburger VC. *Adolescents and the Media: Medical and Psychological Impact*. Thousand Oaks, CA: Sage; 1995
9. Huston AC, Donnerstein E, Fairchild H, et al. *Big World, Small Screen: The Role of Television in American Society*. Lincoln, NE: University of Nebraska Press; 1992

10. Donnerstein E, Linz D. The mass media: a role in injury causation and prevention. *Adolesc Med*. 1995;6:271–284
11. Eron LR. Media violence. *Pediatr Ann*. 1995;24:84–87
12. Willis E, Strasburger VC. Media violence. *Pediatr Clin North Am*. 1998; 45:319–331
13. Kunkel D, Cope KM, Farinola WJM, Biely E, Rollin E, Donnerstein E. *Sex on TV: Content and Context*. Menlo Park, CA: The Henry J Kaiser Family Foundation; 1999
14. Huston AC, Wartella E, Donnerstein E. *Measuring, the Effects of Sexual Content in the Media*. Menlo Park, CA: The Henry J Kaiser Family Foundation Report; 1998
15. Brown JD, Greenberg BS, Buerkel-Rothfuss NL. Mass media, sex and sexuality. *Adolesc Med*. 1993;4:511–525
16. Morgan M. Television and school performance. *Adolesc Med*. 1993;4: 607–622
17. Harrison K, Cantor J. The relationship between media consumption and eating disorders. *J Commun*. 1997;47:40–67
18. Signorelli N. Sex roles and stereotyping on television. *Adolesc Med*. 1993;4:551–561
19. A Different World. *Children's Perceptions of Race and Class in the Media*. Oakland, CA: Children Now; 1998
20. Andersen RE, Crespo CJ, Bartlett SJ, Cheskin LJ, Pratt M. Relationship of physical activity and television watching with body weight and level of fatness among children: results from the Third National Health and Nutrition Examination Study. *JAMA*. 1998;279:938–942
21. Jeffrey RW, French SA. Epidemic obesity in the United States: are fast foods and television viewing contributing? *Am J Public Health*. 1998;88: 277–280
22. Comstock GC, Strasburger VC. Media violence: Q & A. *Adolesc Med*. 1993;4:495–509
23. Federman J, ed. *National Television Violence Study*. Vol 3. Thousand Oaks, CA: Sage; 1998
24. DuRant RH, Rich M, Emans SJ, Rome ES, Allred E, Woods ER. Violence and weapon carrying in music videos: a content analysis. *Arch Pediatr Adolesc Med*. 1997;151:443–448
25. Cantor J. "Mommy, I'm Scared": How TV and Movies Frighten Children and What We Can Do to Protect Them. New York, NY: Harcourt Brace; 1998
26. Cantor J, Nathanson AI. Children's fright reactions to television news. *J Commun*. 1996;46:139–152
27. Gerbner G, Ozyegin N. *Alcohol, Tobacco, and Illicit Drugs in Entertainment Television, Commercials, News, "Reality Shows," Movies, and Music Channels*. Princeton, NJ: Robert Wood Johnson Foundation; 1997
28. DuRant RH, Rome ES, Rich M, Allred E, Emans SJ, Woods ER. Tobacco and alcohol use behaviors portrayed in music videos: a content analysis. *Am J Public Health*. 1997;87:1131–1135
29. Robinson TN, Chen HL, Killen JD. Television and music video exposure and risk of adolescent alcohol use. *Pediatrics* [serial online]. 1998;102:e54. Available at: <http://www.pediatrics.org/cgi/content/full/102/5/e54>. Accessed May 2, 2000.
30. Brown JD, Steele, JR. *Sex and the Mass Media*. Menlo Park, CA: The Henry J Kaiser Family Foundation; 1995
31. Kunkel D, Cope KM, Colvin C. *Sexual Messages on Family Hour Television: Content and Context*. Menlo Park, CA: Henry J Kaiser Family Foundation; 1996
32. Telecommunications Act of., Pub L No. 104–104, 1996.
33. Cantor J. Ratings for program content: the role of research findings. *Ann Am Acad Pol Soc Science*. 1998;557:54–69
34. Kunkel D, Farinola WJM, Cope KM, Donnerstein E et al. *Rating, the TV Ratings: One Year Out. An Assessment of the Television Industry's Use of V-Chip Ratings*. Menlo Park, CA: Henry J Kaiser Family Foundation; 1998
35. *Parents Rate the TV Ratings*. Minneapolis, MN: National Institute on Media and the Family; 1998
36. Potter WJ. *Media Literacy*. Thousand Oaks, CA: Sage; 1998
37. Huesman LR, Eron LD, Klein R, Brice P, Fischer P. Mitigating, the imitation of aggressive behaviors by changing children's attitudes about media violence. *J Pers Soc Psychol*. 1983;44:899–910
38. Gunter B. The question of media violence. In: Bryant J, Zillmann D, eds. *Media Effects: Advances in Theory and Research*. Hillsdale, NJ: Lawrence Erlbaum; 1994:163–211
39. Kubey RW. Television dependence, diagnosis, and prevention. In: MacBeth TM, ed. *Tuning in to Young Viewers: Social Science Perspectives on Television*. Thousand Oaks, CA: Sage; 1996:221–260
40. Singer DG, Singer JL. Developing critical viewing skills and media literacy in children. In: Jordan AB, Jamieson KH, eds. *Children and television*. *Ann Am Acad Pol Soc Science*. 1998;557:164–179

41. Austin EW, Johnson KK. Effects of general and alcohol-specific media literacy training on children's decision making about alcohol. *J Health Commun.* 1997;2:17-42
42. Austin EW, Pinkleton BE, Fujioka Y. The role of interpretation processes and parental discussion in the media's effects on adolescents' use of alcohol. *Pediatrics.* 2000;105:343-349
43. Kubey R, Baker F. Has media literacy found a curricular foothold? *Education Week.* 1999;19:38,56
44. Children's Television Act. 47 USC §303a, 303b, 394
45. Revision of Programming Policies for Television Broadcast Stations. Washington, DC. Federal Communications Commission; August 8, 1996. FCC 96-335 (MM Docket 93-48)
46. American Academy of Pediatrics. *Media Matters: A National Media Education Campaign.* Elk Grove Village, IL: American Academy of Pediatrics; 1997
47. Valerio M, Amodio P, Dal Zio M, Vianello A, Zacchello GP. The use of television in 2- to 8-year-old children and the attitude of parents about such use. *Arch Pediatr Adolesc Med.* 1997;151:22-26
48. TV Turnoff Network Web site. Available at: <http://www.tvturnoff.org>. Accessed December 27, 2000

ADDENDUM

A policy statement on "Developmental Anomalies of the External Genitalia in the Newborn" has recently been published (*Pediatrics.* 2000;106:138-142). The purpose of this review is to identify which newborns among those with abnormal genital development need to be screened for intersexuality, to outline the investigations necessary, and to suggest indications for referral to a center with experience in the diagnosis and management of these disorders.

The 1996 policy on Timing of Elective Surgery states that "children whose genetic sexes are not clearly reflected in external genitalia (ie, hermaphroditism) can be raised successfully as members of either sex if the process begins before the age of 2 years" [see the heading under Body Image and Sexual Development]. The 2000 policy on Developmental Anomalies of the External Genitalia acknowledges the considerable recent debate about the appropriate gender assignment of newborns with the most extreme forms of genital ambiguity, and notes that some have suggested that the current early surgical treatment be abandoned in favor of allowing the affected person to participate in gender assignment at a later time.

This controversy about gender reassignment does not invalidate the other recommendations about the timing of elective surgery on the genitalia of male children with particular reference to the risks, benefits, and psychological effects of surgery and anesthesia that are present in the 1996 statement.